INDEX

Achrestic anaemia, 69.

Agglutinins, and pneumonia, 251; method used to detect, 252; presence of, to homologous antigen in course of pneumonia, 253; absence of, in fatal cases of pneumonia, 254; after the administration of divalent serum, 255.

Anaemia, achrestic, 69.

Anterior pituitary extract on blood-sugar in coeliac disease, effect of, 244; on the blood non-protein nitrogen curve in coeliac disease, effect of, 246; on fat retention and the general condition of the patient with coeliac disease, the effect of, 247.

Asthma, respiratory efficiency tests in, 287; Knipping spirometer results, 288; X-ray screening of the diaphragm, 291; chest measurements, 292; mercury U-tube tests, 293.

Auricular fibrillation, quinidine treatment of, 205.

Bendien-Lowe test, 141.

Beri-beri, chemical and clinical findings in, with special reference to vitamin B_1 deficiency, 355.

Blood, relative sugar content of, and cerebrospinal fluid, 159; relative non-protein nitrogen content of, and cerebrospinal fluid, 162.

Blood lipids, thyroid disease and, 455.

Blood-sugar curve in healthy children, 228; in coeliac disease, 229; flat, not pathognomonic of coeliac disease, 231; flat, cause of, in coeliac disease, 233.

Bronchi, secondary thrush of, 375; distribution of 'Monilia' in Nature, 383; distribution of 'Monilia' in the healthy human respiratory tract, 384; incidence of 'Monilia' in tuberculous sputum, 384; incidence of 'Monilia' in sputum and respiratory tract in non-tuberculous pulmonary disease, 386.

Calcium, role of, in spontaneous overbreathing letany, 341.

Carbohydrate metabolism in coeliac disease, 227.

Cerebrospinal fluid, mechanism of production of, 159; relative sugar content of blood and, 159; relative non-protein nitrogen content of blood and, 162; effect of induced hyperglycaemia on the glucose content of, 169; glucose content of, after withdrawal, 183.

Coeliac disease: *Carbohydrate metabolism*, 227; blood-sugar curve in patients with, 229; flat blood-sugar curve not pathognomonic of, 231; cause of flat blood-sugar curve in, 233; *Effect of insulin and of active pituitary extract*, 242; effect of insulin in, 242; effect of active pituitary extract on blood-sugar in, 244; effect of anterior pituitary extract on the blood non-protein nitrogen curve in, 246; effect of anterior pituitary extract on fat retention and general condition of the patient, 247.

Disinfectants, hypochlorite, killing effect of, on *L. icterohaemorrhagiae*, 280.

Erythrocyte sedimentation rate in diseases of the heart, 1.

Fever therapy, 187.

Gargoylism, 119.

Gaucher's disease, 517.

Gee-Thaysen disease, 481.

Glucose content of cerebrospinal fluid, effect of induced hyperglycaemia, 169; after withdrawal, 183.

Glycogen disease, 31.

Haemo-respiratory exchange, 448.

Haemorrhage, subarachnoid, 461.

Heart, erythrocyte sedimentation rate in diseases of, 1.

Hepato(nephro)megalia glycogenica, 31.

Hyperglycaemia, induced, effect of, on glucose content of cerebrospinal fluid, 169; production of a sustained hyperglycaemia, 170; effect of sustained hyperglycaemia on the C. S. F. sugar content, 172.

Insulin, effect of, in coeliac disease, 242.

Leucocytosis in typed lobar pneumonia: serum treated cases, 105; in patients with positive blood cultures as related

to age, 113; leucocyte count as a prognostic indication, 113; value of leucocyte count in identifying the type, 115.
Lung volume and respiratory exchange after pneumothorax, 445; haemo-respiratory exchange, 448.

'Monilia', distribution of, in Nature, 383; distribution of, in healthy human respiratory tract, 384; incidence of, in tuberculous sputum, 384; incidence of, in sputum and respiratory tract in non-tuberculous pulmonary disease, 386.
Muscular degeneration occurring in late adult life, 51.
Myasthenia gravis, observations on the treatment of, 21; creatine and creatinine metabolism, 23; ephedrine, 23; glycine, 24; parathormone and phosphorus metabolism, 25; acetyl-choline, 26; prostigmin and eserine, 26.

Pneumonia, studies in, with special reference to agglutinins, 251; method used to detect agglutinins, 252; presence of agglutinins to homologous antigen in course of, 253; absence of agglutinins in fatal cases of, 254; serum treatment, 257.
Pneumonia, typed lobar, leucocytosis in, 105; value of leucocyte count in identifying the type, 115.
Pneumothorax, lung volume and respiratory exchange after, 445; haemo-respiratory exchange, 448.
Pneumothorax, artificial, pulmonary congestion following, 327.
Porphyrinuria, congenital, 473.
Pulmonary congestion following artificial pneumothorax, 327.
Pulmonary disease, non-tuberculous, incidence of 'Monilia' in sputum and respiratory tract in, 386.
Pyrexia, therapeutic, by means of short radio waves, 187; development of fever therapy, 187.

Quinidine treatment of auricular fibrillation, 205.

Raynaud phenomena, 399; in normal persons who are exposed to cold long enough to lower the blood temperature, 403; hereditary cold fingers, 405; Raynaud's disease, 408; after local injury to hands or feet, and in workers with vibrating tools, &c., 415; sclerodactyly, 417; thrombo-angeitis obliterans, 422; arterio-sclerosis, 426; syphilitic arteritis, 429; 'rheumatic' (? streptococcal) arteritis, 430; cervical rib, 432; in advanced pulmonary tuberculosis, leukaemia, lupus erythematosus, malaria, polycythaemia vera, chronic arsenical poisoning, &c., 433; sympathetic ganglionectomy, 438.
Respiratory efficiency tests in asthma, 287; Knipping spirometer results, 288; X-ray screening of the movements of the diaphragm, 291; chest measurements, 292; mercury U-tube tests, 293.

Steatorrhoea, idiopathic, of adults and adolescents in non-tropical countries, 481.

Tetany, spontaneous overbreathing, role of calcium in, 341.
Thrush, secondary, of the bronchi, 375; distribution of 'Monilia' in Nature, 383; distribution of 'Monilia' in the healthy human respiratory tract, 384; incidence of 'Monilia' in tuberculous sputum, 384; incidence of 'Monilia' in sputum and respiratory tract in non-tuberculous pulmonary disease, 386.
Thyroid disease and blood lipids, 455.
Tuberculous sputum, incidence of 'Monilia' in, 384.

Vanadate serum malignancy test, modified, 141.
Vitamin B_1 deficiency, chemical and clinical findings in beri-beri with special reference to, 355.
Von Gierke's disease, 31.

Weil's disease in fish-workers, 263; in general population, 283; an occupational disease, 284.
Whooping-cough, review of recent work on, 307; incubation period and mode of infection, 307; length of infectivity, 310; vaccine prophylaxis, 312.